

University of Groningen

Development of an efficient electroporation method for rhizobacterial *Bacillus mycoides* strains

Yi, Yanglei; Kuipers, Oscar P.

Published in:
Journal of microbiological methods

DOI:
[10.1016/j.mimet.2016.12.022](https://doi.org/10.1016/j.mimet.2016.12.022)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Yi, Y., & Kuipers, O. P. (2017). Development of an efficient electroporation method for rhizobacterial *Bacillus mycoides* strains. *Journal of microbiological methods*, 133, 82-86.
<https://doi.org/10.1016/j.mimet.2016.12.022>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Development of an efficient electroporation method for rhizobacterial *Bacillus mycoides* strains

Yanglei Yi, Oscar P. Kuipers *

Molecular Genetics, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Groningen, The Netherlands



ARTICLE INFO

Article history:

Received 3 December 2016

Received in revised form 23 December 2016

Accepted 23 December 2016

Available online 30 December 2016

Keywords:

Bacillus mycoides

Electroporation

Transformation

Optimization

ABSTRACT

In order to develop a method for electroporation of environmental *Bacillus mycoides* strains, we optimized several conditions that affect the electroporation efficiency of this bacterium. By combining the optimized conditions, the electroporation efficiency of strain EC18 was improved to $(1.3 \pm 0.6) \times 10^5$ cfu/ μ g DNA, which is about 10^3 -fold increase in comparison with a previously reported value. The method was further validated on various *B. mycoides* strains, yielding reasonable transformation efficiencies. Furthermore, we confirmed that restriction/modification is the main barrier for electroporation of this bacterium. To the best of our knowledge, this is the first systematic investigation of various parameters of electroporation of *B. mycoides*. The electroporation method reported will allow for efficient genetic manipulation of this bacterium.

© 2017 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Bacillus mycoides is a spore-forming Gram-positive bacterium that is commonly found in soil and the rhizosphere. It belongs to the *B. cereus* sensu lato group, which includes *B. cereus*, *B. thuringiensis*, and *B. anthracis*. *B. mycoides* has received the least attention among this group because it is not a human pathogen as *B. anthracis* and *B. cereus*. Moreover, there are several reports on the insecticidal effect of *B. thuringiensis*, which stimulated further studies using genetic modification (Turchi et al., 2012). The *B. mycoides* species features a unique filamentous growth pattern, either being rotated clockwise or counter-clockwise (Di Franco et al., 2002). Nowadays, more and more studies on *B. mycoides* are focusing on their plant-growth promoting activities (Ambrosini et al., 2016; Bargabus et al., 2002; Neher et al., 2009).

In order to study the plant interaction mechanism of *B. mycoides*, it is necessary to be able to perform molecular genetics studies on this bacterial species. A high transformation efficiency is required to establish the genetic manipulation systems, e.g., enabling gene deletion and mutation within the genome of this organism. Several techniques, including phage transduction, protoplast transformation, natural competence, and electroporation have been applied to incorporate exogenous DNA into *Bacillus* cells (Barlass et al., 2002; Lu et al., 2012; Romero et al., 2006). Among these methods, electroporation is usually the quickest and most reproducible method. Di Franco et al. (2002) modified the electrotransformation protocol of Macaluso and Mettus

(1991) for *B. thuringiensis*, but the implementation on *B. mycoides* resulted in a very low efficiency (less than 200 cfu/ μ g plasmid DNA). Our preliminary experiments show that by applying the method reported by Ehling-Schulz et al. (2005), which was originally developed for *B. cereus*, the *B. mycoides* strain EC18 could be transformed, albeit at a low efficiency. Apart from this, no other protocols have been developed on electroporation methods for *B. mycoides*. This work aimed at developing an efficient electroporation method for several environmental *B. mycoides* strains. *B. mycoides* EC18 which was isolated from the endosphere of potato, and displaying potential plant growth-promoting activity (data not shown), was used as a first testing strain. Factors including growth media, growth phase, electroporation buffer, pulse strength, and incubation time that can affect electroporation efficiency were optimized. As a result, a high electroporation efficiency of $(1.3 \pm 0.6) \times 10^5$ cfu/ μ g DNA was obtained. Furthermore, we evaluated our optimized protocol on other *B. mycoides* strains as well, which resulted in electroporation efficiencies ranging from $(7.3 \pm 2.1) \times 10^2$ to $(1.3 \pm 0.6) \times 10^5$ cfu/ μ g DNA.

2. Materials and methods

2.1. Bacterial strains, plasmids, and media

Escherichia coli strains were grown in Luria broth at 37 °C, 220 rpm. *B. mycoides* strains were isolated from a potato field in Wijster (the Netherlands) (Table 3) and were grown at 30 °C, 200 rpm in LB. When necessary, 100 μ g/ml of ampicillin or 4 μ g/ml of chloramphenicol was added to the culture medium. Plasmid DNA from *E. coli* was purified using the NucleoSpin plasmid isolation kit (Macherey-Nagel GmbH & Co. Düren, Germany) according to the manufacturer's instructions. A

* Corresponding author at: Molecular Genetics Group, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Nijenborgh 7, 9747 AG Groningen, The Netherlands.

E-mail address: o.p.kuipers@rug.nl (O.P. Kuipers).

Geobacillus-E. coli shuttle vector pNW33N (Bacillus Genetic Stock Centre) isolated from *E. coli* MC1061 was used as the plasmid for electroporation protocol optimization. pNW33N isolated from *E. coli* JM110 was used to test the plasmid methylation effects on transformation efficiency.

2.2. Screening for growth medium

The *B. mycoides* strain was streaked on LB agar plate and grown at 30 °C overnight. One single colony was inoculated into various media including LB (10 g tryptone, 5 g yeast extract, 10 g NaCl in 1 L deionized water, pH 7.2), LBS (10 g tryptone, 5 g yeast extract, 10 g NaCl, and 91.1 g sorbitol in 1 L deionized water, pH 7.2), LBSP (10 g tryptone, 5 g yeast extract, 10 g NaCl, 50 mM KH₂PO₄ and K₂HPO₄, and 91.1 g sorbitol in 1 L deionized water, pH 7.2), 2×YT (16 g tryptone, 10 g yeast extract and 5 g NaCl in 1 L deionized water, pH 7.2), and BHIS (34 g BHI, and 91.1 g sorbitol in 1 L deionized water, pH 7.2).

2.3. Preparation of electro-competent cells

B. mycoides cells were grown overnight in an appropriate medium at 30 °C, 200 rpm. The culture was diluted with the same medium to obtain an initial optical density at 600 nm (OD₆₀₀) of about 0.05 then continually grown. The OD₆₀₀ was measured by a Genesys 20 spectrophotometer (ThermoSpectronic, USA) during bacterial growth. When the OD₆₀₀ reached the appropriate value, the cell culture was transferred into a 50 mL centrifuge tube and cooled on ice for 10 min. Cells were then collected by centrifugation at 4 °C, 3000 ×g for 10 min. After being washed four times in corresponding electroporation buffer (pre-chilled), the cell pellets were re-suspended in 1 mL of the electroporation buffer. The resulting electro-competent cells were flash-frozen in liquid nitrogen and stored at −80 °C prior to electroporation.

2.4. Cell wall-weakening treatment

Serial concentrations of Glycine or DL-threonine were added when bacterial cultures reached an OD₆₀₀ = 0.85 for the cell-wall weakening treatment. These treated bacterial cultures were continued to shake for 1 h. After that, electro-competent cells were prepared with the methods mentioned above.

2.5. Electroporation

The electroporation was performed as previously reported method (Peng et al., 2009; Turgeon et al., 2006) with slight modification. 100 µL of frozen electro-competent cells was thawed on ice and mixed with 2–3 µL of plasmid DNA (1 µg). Cells mixed with the same amount of deionized water served as negative control. The mixture was loaded into an electroporation cuvette (2 mm electrode gap, pre-chilled) and exposed to a single pulse in a Gene Pulser System (Bio-Rad, USA) with the settings of 25 µF capacitance, 100, 200, or 400 Ω resistance, and voltage ranges between 5 and 12.5 kV cm^{−1}. After pulse-shock, cells were immediately added with 1 mL of pre-warmed corresponding growth medium and transferred to a 2 mL Eppendorf tube. After 2 h incubation at 30 °C, 180 rpm, the dilutions of the recovered cell culture were plated on LB agar with 4 µg/mL of chloramphenicol. The plates were incubated at 30 °C overnight and transformation efficiencies (cfu/µg DNA) were calculated by counting the colonies on plates.

3. Results and discussion

3.1. Optimization of growth conditions

For determining the optimal growth medium, we transformed *B. mycoides* according to the protocol published by Ehling-Schulz et al.

(2005). Strain EC18 was grown in different media before preparing the electro-competent cells. Media including LB, LBS, LBSP, 2×YT and BHIS were chosen according to reported electroporation methods for *Bacillus* species (Zhang et al., 2011a, 2015). When the OD₆₀₀ reached a value of 0.6, the competent cells were prepared by washing the cell pellet with increasing concentrations of ice-cold glycerol (2.5, 5, and 10%). Electroporation was performed with 1 µg pNW33N plasmid DNA and recovered for 2 h with the corresponding growth media. The efficiencies were calculated after 1 day (Table 1) on 4 µg/mL chloramphenicol selection media. Among the five tested media, the super rich medium BHIS showed the highest efficiency of $(6.2 \pm 1.4) \times 10^3$ cfu/µg plasmid DNA. Zhang et al. (2011a) reported that the transformation efficiency of *B. amyloliquefaciens* was positively correlated to concentrations of salts, whereas negatively related to the nutritional ingredient concentration. Surprisingly, we observed an opposite effect of the nutritional and salt concentrations on transformation efficiencies in our study. The transformation efficiency by culturing in the LBSP medium is lower than culturing in LBS medium. Moreover, *B. mycoides* EC18 could not grow in the hypertonic media NCM and M9YE (data not shown). We hypothesized that *B. mycoides* EC18 is sensitive to salt concentrations.

According to previous reports, cells of *B. cereus* and *B. thuringiensis* collected at early growth-stage show better electroporation efficiency than late growth-stage cultures (Peng et al., 2009; Turgeon et al., 2006), while *B. subtilis* WB800 has high electroporation efficiency of 1.88×10^5 cfu/µg DNA at late growth-stage (OD₆₀₀ = 2.2–2.3) (Lu et al., 2012). To investigate the effects of the growth phase of *B. mycoides* on the electroporation efficiency, *B. mycoides* cells were cultured in BHIS medium to OD₆₀₀ from 0.3 to 1.8 for competent cell preparations. Our results indicated that when OD₆₀₀ is between 0.9 and 1, the highest electroporation efficiency was obtained (Fig. 1A). According to the growth curve (Fig. 1B), cells are in the early stage of exponential growth. A similar phenomenon was also reported for *B. subtilis* ZK by Zhang et al. (2015). This indicates that cells from the early exponential phase result in higher electroporation efficiency of *B. mycoides*.

3.2. Optimization of cell wall-weakening agents

The cell wall-weakening agents, glycine and DL-threonine, are widely used to improve the electroporation efficiency of environmental strains. These amino acids can reduce the peptidoglycan bonds and loosen up the cell wall by replacing the L- and D-alanine bridges (Hammes et al., 1973). In this study, EC18 cells were first grown in BHIS medium, and when the OD₆₀₀ reached about 0.85, glycine and threonine were added at different concentrations (0%, 1%, 2%, 3%, 4% and 5%). After 1 h of additional incubation, the cells were collected and concentrated to obtain competent cells. Electroporation treatments were performed and the greatest transformation efficiency was obtained in the 2% glycine treatment group. The same concentration of threonine also improved the transformation efficiency, which was however lower than that of glycine (Fig. 2). Cell growth rate was slightly reduced in the presence of both glycine and DL-threonine, and high concentrations of glycine treatments resulted in cellular lysis in the samples

Table 1
Effect of growth media on the electroporation efficiency of *B. mycoides* EC18.

Medium	Transformation efficiency (cfu/µg DNA)
LB	$(3.4 \pm 1.0) \times 10^2$
LBS	$(8.3 \pm 2.5) \times 10^2$
LBSP	$(2.0 \pm 1.1) \times 10^2$
2×YT	$(5.4 \pm 1.3) \times 10^2$
BHIS	$(6.2 \pm 1.4) \times 10^3$

Cells were grown in different media to OD₆₀₀ ~0.6 to prepare the electro-competent cells using series concentration of glycerol solution as electroporation buffer. 1 µg of pNW33N plasmid was used for electroporation with the settings 25 µF, 10 kV cm^{−1}, 200 Ω. Data are shown as mean ± standard deviation based on 3 replications.




Fig. 1. Effect of growth phase on the electroporation efficiency of *B. mycoides* EC18. (A) The transformation efficiency at different growth phases of *B. mycoides* EC18. Cells were grown in BHIS medium to various OD₆₀₀ densities for the electro-competent cell preparation, using serial concentrations of a glycerol solution as electroporation buffer. 1 µg of pNW33N plasmid was used for electroporation with the settings 25 µF, 10 kV cm⁻¹, 200 Ω. (B) The growth of *B. mycoides* EC18 in BHIS medium at 30 °C, 200 rpm. At different time points, the optical density of the cell cultures was measured spectrophotometrically (OD_{600 nm}) (N = 3, bar = standard deviation).

(data not shown). These results demonstrate that the transformation efficiency of *B. mycoides* has been significantly enhanced by a cell wall-weakening treatment. The optimal results were obtained at incubations with 2% glycine.

3.3. Effects of buffer on the electroporation efficiency

According to previous studies, the composition of the electroporation buffer has a tremendous impact on transformation efficiency. Xue et al. (1998) improved the transformation efficiency of *B. subtilis* and *B. licheniformis* by applying electroporation buffers with high osmolality. It has also been reported that with the addition of trehalose, the transformation efficiency of *B. subtilis* can be enhanced dramatically to about 100-fold (Lu et al., 2012). In this study, EC18 cells growing in BHIS medium was added with 2% glycine when the cell density reached an OD₆₀₀ = 0.85. After 1 h of additional incubation, cells were collected (OD₆₀₀ ~0.95) and washed four times in different electroporation buffers (Table 2) to obtain competent cells. The transformation

efficiency is shown in Table 2. The buffers consisting of glycerol exhibited better transformation efficiencies than those with sucrose. The highest transformation efficiency was obtained with buffer G containing 10% glycerol, 0.25 M sorbitol, and 0.25 M trehalose. However, fewer transformants were obtained when the sorbitol and trehalose concentration increased to 0.5 M. The addition of salts (buffers K, L, and M) showed no improvement on transformation efficiency. Based on these results, buffer G was used throughout.

3.4. Effects of the electric field on the electroporation efficiency

A relatively high electrical field is necessary to create pores in the cells and thus provide a temporary pathway for exogenous DNA. After the electric pulse, they gradually reseal and most cells recover. However, if the electrical field is above a certain level, most pores either do not reseal or reseal too slowly to preserve cell viability (Kotnik et al., 2015). To optimize the electric field for *B. mycoides* EC18, a gradient of field strength (7.5–12.5 kV cm⁻¹) and resistance values (100, 200, and 400 Ω) at 25 µF was tested. The results are shown in Fig. 3, an electric field of 10 kV cm⁻¹ and resistance of 200 Ω led to the optimal transformation efficiency. This electric field is lower than that of *B. subtilis* (Lu et al., 2012; Zhang et al., 2015) and closely related species, e.g., *B. cereus* and *B. thuringiensis* (Peng et al., 2009; Turgeon et al., 2006). These results indicate that *B. mycoides* is more vulnerable to high electric fields compared to other *Bacillus* species.

3.5. Effects of recovery time

After an electric pulse, the injured bacterial cells need several hours to retain viability and to allow phenotypic expression. To optimize the recovery time, the pulse-shocked cells were immediately transferred to BHIS medium and incubated at 30 °C, 180 rpm for 2–7 h. As shown in Fig. 4, the transformation efficiency was increased as the incubation time extended. Remarkably, the transformation efficiency reached 1.2 × 10⁵ cfu/µg DNA after 5 h of incubation. Further elongation of the incubation time only led to a slight increase in transformation efficiency. Hence, we used 5 h of recovery time for *B. mycoides*. Xue et al. (1998) pointed out that the increased transformation efficiency at the longer recovery times might be due to the division of transformed cells. Zhang et al. (2011b) also used a long recovery time of 5 hrs for electroporation of *Arthrobacter*.




Fig. 2. Effect of wall-weakening agents on the electroporation efficiency of *B. mycoides* EC18. Cells were grown in BHIS medium to OD₆₀₀ ~0.85, then incubated 1 h with different concentrations of cell wall-weakening agents. Electro-competent cells were prepared using glycerol solutions as electroporation buffer. 1 µg of pNW33N plasmid was used for electroporation with the settings 25 µF, 10 kV cm⁻¹, 200 Ω (N = 3, bar = standard deviation).

Table 2Effect of electroporation buffers on transformation efficiencies in *B. mycoides* EC18.

Buffer	Base component	Salt component	Transformation efficiency (cfu/ μ g DNA)	Reference
A	Glycerol solution (2.5, 5, and 10%)	–	$(4.6 \pm 1.7) \times 10^4$	Ehling-Schulz et al. (2005)
B	10% glycerol	–	$(4.0 \pm 2.0) \times 10^4$	Zhang et al. (2011b)
C	10% glycerol, 0.25 M sorbitol	–	$(6.2 \pm 2.8) \times 10^4$	Zhang et al. (2011b)
D	10% glycerol, 0.5 M sorbitol	–	$(5.8 \pm 1.8) \times 10^4$	Zhang et al. (2011b)
E	10% glycerol, 0.25 M sorbitol, 0.25 M mannitol	–	$(4.2 \pm 1.7) \times 10^4$	This study
F	10% glycerol, 0.5 M sorbitol, 0.5 M mannitol	–	$(4.0 \pm 0.8) \times 10^2$	Xue et al. (1998)
G	10% glycerol, 0.25 M sorbitol, 0.25 M trehalose	–	$(6.8 \pm 1.6) \times 10^4$	This study
H	10% glycerol, 0.5 M sorbitol, 0.5 M trehalose	–	$(4.0 \pm 0.8) \times 10^2$	This study
I	10% glycerol, 0.25 M sorbitol, 0.25 M mannitol, 0.25 M trehalose	–	$(1.2 \pm 0.8) \times 10^2$	This study
J	0.5 M sorbitol, 0.5 M mannitol, 0.5 M trehalose	–	44 \pm 22	Zhang et al. (2015)
K	0.5 M sorbitol, 0.5 M mannitol, 0.5 M trehalose	0.5 mM MgCl ₂ , 0.5 mM K ₂ HPO ₄ /KH ₂ PO ₄	No transformants	Zhang et al. (2015)
L	272 mM sucrose	0.5 mM MgCl ₂ , 0.5 mM K ₂ HPO ₄ /KH ₂ PO ₄	$(5.2 \pm 2.0) \times 10^2$	Silo-Suh et al. (1994)
M	250 mM sucrose, 10% glycerol	1 mM HEPES; 1 mM MgCl ₂	No transformants	Turgeon et al. (2006)

Cells were grown in BHIS medium to OD₆₀₀ ~0.85, then incubated 1 h with 2% glycine. Electro-competent cells were prepared using different electroporation buffers. 1 μ g of pNW33N plasmid was used for electroporation with the settings 25 μ F, 10 kV cm⁻¹, 200 Ω . Data are shown as mean \pm standard deviation of 3 replications.

3.6. Application of the optimized transformation protocol on other *B. mycoides* strains and the effects of plasmid methylation

In order to validate the optimized protocol in other environmental *B. mycoides* isolates, we combined all the improved factors and applied the method on 5 other strains in addition to strain EC18. Since the restriction of methylated DNA in several *Bacillus* species is known as a major barrier to transformation and genetic manipulation (Sitaraman and Leppla, 2012), we also compared the transformation efficiency of plasmids that were isolated from *E. coli* methylation strain MC1061 (dam⁺/dcm⁺) and methylation-deficient strain JM110 (dam⁻/dcm⁻). As shown in Table 3, for the plasmid isolated from *E. coli* MC1061, only strain EC18 showed high transformation efficiency of $(1.1 \pm 0.6) \times 10^5$ cfu/ μ g DNA. Notably, strain SB4 was transformed at a much lower level of efficiency and the other strains were not transformable at all. However, when non-methylated plasmid DNA was used, this

protocol can efficiently be applied for all the tested *B. mycoides* strains. The transformation efficiencies are ranging from $(7.3 \pm 2.1) \times 10^2$ to $(1.3 \pm 0.6) \times 10^5$ cfu/ μ g DNA. These results indicate that our optimized protocol can be widely used in *B. mycoides* strains. Methylation of transmitted DNA can clearly affect the efficiency of transformation, with the exception of strain EC18, which only showed a slight difference between methylated and non-methylated DNA.

4. Conclusion

This paper describes an efficient electroporation method for several environmental *B. mycoides* strains, which is reproducible and convenient. By using this method, we obtained transformation efficiency up to $(1.3 \pm 0.6) \times 10^5$ cfu/ μ g DNA. We confirmed that our protocol can be applied to several *B. mycoides* strains. The method described here facilitates advanced molecular genetics studies in this important biocontrol bacterium.




Fig. 3. Effect of electric field on the electroporation efficiency of *B. mycoides* EC18. Cells were grown in BHIS medium to OD₆₀₀ ~0.85, then incubated 1 h with 2% glycine. Electro-competent cells were prepared using buffer G. 1 μ g of pNW33N plasmid was used for electroporation with a gradient of field strength (7.5–12.5 kV cm⁻¹) and resistance values (100, 200, and 400 Ω) at 25 μ F (N = 3, bar = standard deviation).




Fig. 4. Effect of incubation time on the electroporation efficiency of *B. mycoides* EC18. Cells were grown in BHIS medium to OD₆₀₀ ~0.85, then incubated 1 h with 2% glycine. Electro-competent cells were prepared using buffer G. 1 μ g of pNW33N plasmid was used for electroporation with the settings 25 μ F, 10 kV cm⁻¹, 200 Ω . Cells were then incubated at 30 °C, 180 rpm for 2–7 h (N = 3, bar = standard deviation).

Table 3

Electroporation of wild type *B. mycoides* strains with methylated and non-methylated plasmid DNA.

Strain	Electroporation efficiency of different plasmid source (cfu/μg DNA)	
	<i>E. coli</i> MC 1061 (dam ⁺ /dcm ⁺)	<i>E. coli</i> JM110 (dam [−] /dcm [−])
<i>B. mycoides</i> EC18	$(1.1 \pm 0.6) \times 10^5$	$(1.3 \pm 0.6) \times 10^5$
<i>B. mycoides</i> M2E15	–	$(3.7 \pm 1.2) \times 10^3$
<i>B. mycoides</i> S2E19	–	$(1.7 \pm 0.4) \times 10^4$
<i>B. mycoides</i> S3E15	–	$(7.3 \pm 2.1) \times 10^2$
<i>B. mycoides</i> SB4	$(1.8 \pm 0.6) \times 10^3$	$(6.4 \pm 1.2) \times 10^4$
<i>B. mycoides</i> SB8	–	$(5.7 \pm 1.6) \times 10^4$

Cells were grown in BHIS medium to OD₆₀₀ ~0.85, then incubated 1 h with 2% glycine. Electro-competent cells were prepared using buffer G. 1 μg of PNW33N plasmid were used for electroporation with the settings as: 25 μF, 10 kV cm^{−1}, 200 Ω. Cells were then incubated at 30 °C, 180 rpm for 5 h. Data are shown as mean ± standard deviation of 3 replications.

Acknowledgement

We are grateful to Prof. J.D. van Elsas and J. Spoelder (University of Groningen, the Netherlands) for providing the *B. mycoides* strains. We thank Dr. C. Song in our group for critical reading and suggestions. Y. Yi was supported by a scholarship from the China Scholarship Council (201306300040).

References

- Ambrosini, A., Stefanski, T., Lisboa, B.B., Beneduzi, A., Vargas, L.K., Passaglia, L.M.P., 2016. Diazotrophic bacilli isolated from the sunflower rhizosphere and the potential of *Bacillus mycoides* B38V as biofertiliser. *Ann. Appl. Biol.* 168, 93–110.
- Bargabus, R., Zidack, N., Sherwood, J., Jacobsen, B., 2002. Characterisation of systemic resistance in sugar beet elicited by a non-pathogenic, phyllosphere-colonizing *Bacillus mycoides*, biological control agent. *Physiol. Mol. Plant Pathol.* 61, 289–298.
- Barlass, P.J., Houston, C.W., Clements, M.O., Moir, A., 2002. Germination of *Bacillus cereus* spores in response to L-alanine and to inosine: the roles of *gerL* and *gerQ* operons. *Microbiology* 148, 2089–2095.
- Di Franco, C., Beccari, E., Santini, T., Pisaneschi, G., Tecce, G., 2002. Colony shape as a genetic trait in the pattern-forming *Bacillus mycoides*. *BMC Microbiol.* 2, 33.

- Ehling-Schulz, M., Vukov, N., Schulz, A., Shaheen, R., Andersson, M., Märklbauer, E., et al., 2005. Identification and partial characterization of the nonribosomal peptide synthetase gene responsible for cereulide production in emetic *Bacillus cereus*. *Appl. Environ. Microbiol.* 71, 105–113.
- Hammes, W., Schleifer, K., Kandler, O., 1973. Mode of action of glycine on the biosynthesis of peptidoglycan. *J. Bacteriol.* 116, 1029–1053.
- Kotnik, T., Frey, W., Sack, M., Haberl Meglic, S., Peterka, M., Miklavcic, D., 2015. Electroporation-based applications in biotechnology. *Trends Biotechnol.* 33, 480–488.
- Lu, Y.P., Zhang, C., Lv, F.X., Bie, X.M., Lu, Z.X., 2012. Study on the electro-transformation conditions of improving transformation efficiency for *Bacillus subtilis*. *Lett. Appl. Microbiol.* 55, 9–14.
- Macaluso, A., Mettus, A.M., 1991. Efficient transformation of *Bacillus thuringiensis* requires nonmethylated plasmid DNA. *J. Bacteriol.* 173, 1353–1356.
- Neher, O.T., Johnston, M.R., Zidack, N.K., Jacobsen, B.J., 2009. Evaluation of *Bacillus mycoides* isolate BmJ and *B. mojavensis* isolate 203-7 for the control of anthracnose of cucurbits caused by *Glomerella cingulata* var. *orbicula*. *Biol. Control* 48, 140–146.
- Peng, D., Luo, Y., Guo, S., Zeng, H., Ju, S., Yu, Z., et al., 2009. Elaboration of an electroporation protocol for large plasmids and wild-type strains of *Bacillus thuringiensis*. *J. Appl. Microbiol.* 106, 1849–1858.
- Romero, D., Perez-Garcia, A., Veening, J.W., de Vicente, A., Kuipers, O.P., 2006. Transformation of undomesticated strains of *Bacillus subtilis* by protoplast electroporation. *J. Microbiol. Methods* 66, 556–559.
- Silo-Suh, L.A., Lethbridge, B.J., Raffel, S.J., He, H., Clardy, J., Handelsman, J., 1994. Biological activities of two fungistatic antibiotics produced by *Bacillus cereus* UW85. *Appl. Environ. Microbiol.* 60, 2023–2030.
- Sitaraman, R., Leppla, S.H., 2012. Methylation-dependent DNA restriction in *Bacillus anthracis*. *Gene* 494, 44–50.
- Turchi, L., Santini, T., Beccari, E., Di Franco, C., 2012. Localization of new peptidoglycan at poles in *Bacillus mycoides*, a member of the *Bacillus cereus* group. *Arch. Microbiol.* 194, 887–892.
- Turgeon, N., Laflamme, C., Ho, J., Duchaine, C., 2006. Elaboration of an electroporation protocol for *Bacillus cereus* ATCC 14579. *J. Microbiol. Methods* 67, 543–548.
- Xue, G.-P., Johnson, J.S., Dalrymple, B.P., 1998. High osmolarity improves the electrotransformation efficiency of the gram-positive bacteria *Bacillus subtilis* and *Bacillus licheniformis*. *J. Microbiol. Methods* 34, 183–191.
- Zhang, G.Q., Bao, P., Zhang, Y., Deng, A.H., Chen, N., Wen, T.Y., 2011a. Enhancing electrotransformation competency of recalcitrant *Bacillus amyloliquefaciens* by combining cell-wall weakening and cell-membrane fluidity disturbing. *Anal. Biochem.* 409, 130–137.
- Zhang, H., Li, Y., Chen, X., Sheng, H., An, L., 2011b. Optimization of electroporation conditions for *Arthrobacter* with plasmid PART2. *J. Microbiol. Methods* 84, 114–120.
- Zhang, Z., Ding, Z.T., Shu, D., Luo, D., Tan, H., 2015. Development of an efficient electroporation method for iturin A-producing *Bacillus subtilis* ZK. *Int. J. Mol. Sci.* 16, 7334–7351.